

Socioeconomic and Environmental Risk Factors for Malaria in Young Children: A Review

Katu Amina H.

School of Natural and Applied Sciences Kampala International Uganda

ABSTRACT

Malaria remained a significant public health challenge in sub-Saharan Africa, disproportionately affecting children under five years of age. Understanding the socioeconomic and environmental risk factors associated with malaria in this vulnerable population was crucial for developing targeted interventions to reduce transmission and improve health outcomes. This review examined the complex interplay between socioeconomic status, housing conditions, environmental factors, and malaria risk among young children. A comprehensive literature search was conducted to synthesize current research findings on the topic. Key socioeconomic determinants included household income, maternal education, and access to preventive measures. Children from low-income families were at higher risk due to limited resources for malaria prevention and treatment, as well as poorer living conditions that facilitate mosquito breeding. Environmental factors such as proximity to stagnant water bodies, inadequate sanitation, and climate variability further exacerbated malaria transmission. The review highlighted the need for multifaceted approaches that address both biological and social determinants of health to effectively reduce the burden of malaria among children under five in sub-Saharan Africa.

Keywords: Malaria, Socioeconomic factors, Environmental factors, young children, Sub-Saharan Africa

INTRODUCTION

Malaria continues to be a leading cause of morbidity and mortality in young children, particularly in sub-Saharan Africa (SSA), where the disease disproportionately affects those under five years of age [1,2]. Despite global efforts to control malaria, significant disparities persist, largely driven by a complex interplay of socioeconomic and environmental factors [3,4]. Understanding these risk factors is crucial for developing effective interventions aimed at reducing malaria transmission and improving health outcomes among vulnerable populations [5]. Socioeconomic factors, including household income, maternal education, and living conditions, have been shown to significantly influence the risk of malaria infection in young children [6]. For instance, children from low-income households or those whose mothers have limited educational attainment are at a higher risk of contracting malaria. These socioeconomic determinants not only affect access to preventive measures such as insecticide-treated nets (ITNs) and healthcare services but also shape community

awareness and practices related to malaria prevention [7,8]. Environmental conditions further exacerbate the risk of malaria among young children. Factors such as proximity to stagnant water bodies, poor housing quality, and inadequate sanitation create favorable breeding grounds for anopheles mosquitoes, the primary vectors of malaria [9]. Rural areas often face greater challenges due to limited infrastructure and resources, increasing the vulnerability of children residing in these settings [10,11]. This review aims to synthesize current knowledge on the socioeconomic and environmental risk factors associated with malaria in young children. By examining existing literature and highlighting key findings, this review seeks to inform public health strategies that address these determinants holistically. Ultimately, understanding and mitigating these risk factors is essential for reducing the burden of malaria in young children and achieving broader health equity goals in sub-Saharan Africa.

SOCIOECONOMIC FACTORS

Socioeconomic factors play a pivotal role in shaping the prevalence and severity of malaria among young children, particularly in sub-Saharan Africa (SSA), where the disease burden is disproportionately high [12]. The interplay between household income, maternal education, living conditions, and access to healthcare significantly influences malaria risk, making it essential to understand these dynamics for effective public health interventions.

- i. **Household Income and Wealth:** One of the most critical socioeconomic determinants of malaria risk is household income or wealth status [13]. Studies have consistently shown that children from low-income households are at a significantly higher risk of contracting malaria compared to those from wealthier families [14]. For instance, a recent analysis revealed that children living in the poorest households had a malaria prevalence rate of 38.2%, while those from the richest households had a prevalence of only 3.6%. Wealthier households have better access to preventive measures like ITNs and IRS, while poorer households lack these resources, increasing their vulnerability to malaria transmission [15,16]. Additionally, higher-income families have better healthcare access, preventing delays and enhancing disease severity.
- ii. **Maternal Education:** Maternal education is another significant factor influencing malaria risk among young children [17]. Research indicates that children whose mothers have secondary education are about 56% less likely to contract malaria compared to those whose mothers have no formal education. Educated mothers are better equipped to understand malaria prevention guidelines and navigate healthcare systems, and their higher education often correlates with increased household income, enhancing access to resources for effective treatment [18,19].
- iii. **Living Conditions:** The quality of living conditions significantly impacts malaria transmission dynamics [20]. Poor housing structures, characterized by inadequate sanitation and proximity to mosquito breeding sites, exacerbate the risk of infection among young children [21]. Housing quality, environmental sanitation, and waste management practices play crucial roles in reducing mosquito entry points and breeding sites, while inadequate

sanitation infrastructure leads to higher malaria transmission rates due to increased vector populations [22].

- iv. **Rural vs. Urban Disparities:** Geographical disparities also play a significant role in malaria risk associated with socioeconomic factors. Rural children face higher risk due to infrastructure limitations, lack of quality healthcare, and agricultural practices that increase mosquito breeding habitats, exacerbate vulnerability and complicate malaria control efforts [23,24].

ENVIRONMENTAL FACTORS

Environmental factors significantly influence the transmission dynamics of malaria, particularly among vulnerable populations such as young children in sub-Saharan Africa. Understanding these factors is crucial for developing effective control strategies and mitigating the impact of malaria [25]. The interplay between natural and human-made environmental conditions, including climatic variables, land use, and sanitation practices, shapes the habitat and behavior of malaria vectors, primarily anopheles mosquitoes [26].

- i. **Climatic Influences:** Malaria transmission is influenced by climatic factors such as temperature, rainfall, and humidity [27]. Temperature is crucial for mosquito development, with optimal temperatures between 20°C and 30°C. Rainfall patterns affect mosquito breeding habitats, with heavy rains leading to increased breeding sites [28]. Humidity is vital for mosquito survival, with higher levels facilitating longer lifespans. Low humidity levels decrease mosquito lifespan, reducing transmission rates. Overall, climatic factors play a significant role in malaria transmission [29].
- ii. **Land Use and Human Activities:** Human activities and land use changes significantly impact malaria transmission dynamics. Agricultural practices, such as irrigated agriculture, can create breeding habitats for mosquitoes, increasing vector densities. Urbanization, on the other hand, can either exacerbate or mitigate malaria transmission, with informal settlements lacking adequate infrastructure and households near stagnant water bodies at greater risk [30-32].
- iii. **Sanitation and Waste Management:** Poor waste management and community awareness are key to controlling malaria transmission [33]. Poor waste management creates mosquito breeding

<https://www.inosr.net/inosr-experimental-sciences/> sites, increasing malaria incidence [34]. Engaging communities in environmental sanitation, such as regular cleaning and

Katu

proper waste disposal, reduces mosquito breeding sites and empowers them to take proactive measures [35-37].

CONCLUSION

Addressing the socioeconomic and environmental determinants of malaria is vital for reducing its burden among children under five in sub-Saharan Africa. Comprehensive strategies that improve household wealth, enhance maternal education, and increase access to healthcare are essential components of effective malaria control programs. Simultaneously, interventions targeting

environmental factors, such as habitat modification and community education on reducing standing water, can significantly contribute to lowering malaria transmission rates. By addressing these multifaceted risk factors, public health initiatives can more effectively protect vulnerable populations from the devastating impacts of malaria.

REFERENCES

1. Sarfo JO, Amoada M, Kordorwu PY, Adams AK, Gyan TB, Osman AG, Asiedu I, Ansah EW. Malaria amongst children under five in sub-Saharan Africa: a scoping review of prevalence, risk factors and preventive interventions. *Eur J Med Res.* 2023 Feb 17;28(1):80. doi: 10.1186/s40001-023-01046-1. PMID: 36800986; PMCID: PMC9936673.
2. Mbishi, J.V., Chombo, S., Luoga, P. *et al.* Malaria in under-five children: prevalence and multi-factor analysis of high-risk African countries. *BMC Public Health* **24**, 1687 (2024). <https://doi.org/10.1186/s12889-024-19206-1>
3. Oshagbemi, O.A., Lopez-Romero, P., Winnips, C. *et al.* Estimated distribution of malaria cases among children in sub-Saharan Africa by specified age categories using data from the Global Burden of Diseases 2019. *Malar J* **22**, 371 (2023). <https://doi.org/10.1186/s12936-023-04811-z>
4. Tsegaye AT, Ayele A, Birhanu S (2021) Prevalence and associated factors of malaria in children under the age of five years in Wogera district, northwest Ethiopia: A cross-sectional study. *PLoS ONE* 16(10): e0257944. <https://doi.org/10.1371/journal.pone.0257944>
5. Coker, M., Folayan, M.O., Michelow, I.C. *et al.* Things must not fall apart: the ripple effects of the COVID-19 pandemic on children in sub-Saharan Africa. *Pediatr Res* **89**, 1078–1086 (2021). <https://doi.org/10.1038/s41390-020-01174-y>
6. Anyanwu PE, Fulton J, Evans E, Paget T. Exploring the role of socioeconomic factors in the development and spread of anti-malarial drug resistance: a qualitative study. *Malar J.* 2017 May 18;16(1):203. doi: 10.1186/s12936-017-1849-1. PMID: 28521791; PMCID: PMC5437569.
7. Ekpono, E. U., Aja, P. M., Ibiham, U. A., Alum, E. U., & Ekpono, U. E. Ethanol Root-extract of *Sphenocentrum jollyanum* Restored Altered Haematological Markers in *Plasmodium berghei*-infected Mice. *Earthline Journal of Chemical Sciences.* 2019; 2(2):189-203. <https://doi.org/10.34198/ejcs.2219.189203>
8. Egwu, C. O., Alope, C., Chukwu, J., Agwu, A., Alum, E., Tsamesidis, I, et al. A world free of malaria: It is time for Africa to actively champion and take leadership of elimination and eradication strategies. *Afr Health Sci.* 2022 Dec;22(4):627-640. doi: 10.4314/ahs.v22i4.68.
9. Mauricio Santos-Vega, Rachel Lowe, Luc Anselin, Vikas Desai, Keshav G Vaishnav, Ashish Naik, Mercedes Pascual, Quantifying climatic and socioeconomic drivers of urban malaria in Surat, India: a statistical spatiotemporal modelling study, *The Lancet Planetary Health*, 2023; 7(12): e985-e998. [https://doi.org/10.1016/S2542-5196\(23\)00249-8](https://doi.org/10.1016/S2542-5196(23)00249-8).
10. Egwu, C.O., Alope, C., Chukwu, J., Nwankwo, J.C., Irem, C., Nwagu, K.E., Nwite, F., Agwu, A.O., Alum, E., Offor, C.E. and Obasi, N.A. Assessment of the Antimalarial Treatment Failure in Ebonyi State, Southeast Nigeria. *J Xenobiot.* 2023 Jan 3;13(1):16-26. doi: 10.3390/jox13010003.
11. Beloconi, A., Nyawanda, B.O., Bigogo, G. *et al.* Malaria, climate variability, and interventions: modelling transmission

- dynamics. *Sci Rep* **13**, 7367 (2023). <https://doi.org/10.1038/s41598-023-33868-8>
12. Kungu, E., Inyangat, R., Ugwu, O.P.C. and Alum, E. U. Exploration of Medicinal Plants Used in the Management of Malaria in Uganda. *Newport International Journal of Research in Medical Sciences*, 2023; 4(1):101-108.<https://nijournals.org/wp-content/uploads/2023/10/NIJ RMS-41101-108-2023.docx.pdf>
13. Obeagu, E. I., Alum, E. U. and Ugwu, O. P. C. Hepcidin's Antimalarial Arsenal: Safeguarding the Host. *Newport International Journal of Public Health and Pharmacy*.2023;4(2):1-8. <https://doi.org/10.59298/NIJPP/2023/10.1.1100>
14. 14 Xing, SY., Zhang, HT., Wang, LM. *et al.* Examining the paradox: increased malaria risk in children under 5 in female-headed households in Nigeria. *Malar J* **23**, 171(2024).<https://doi.org/10.1186/s12936-024-04997-w>
15. Katushabe, J., Nyanzi, J.B. & Muwanga, G.S. Exploring the role of spending on malaria incidence in Uganda using the autoregressive distributed lag approach. *Malar J* **23**,129(2024). <https://doi.org/10.1186/s12936-024-04929-8>
16. 16 Obeagu, E. I., Alum, E. U. and Ugwu, O. P. C. Hepcidin: The Gatekeeper of Iron in Malaria Resistance **NEWPORT INTERNATIONAL JOURNAL OF RESEARCH IN MEDICAL SCIENCES**. 2023;4(2):1-8. <https://doi.org/10.59298/NIJ RMS/2023/10.1.1400>
17. Ritesh Ranjha, Kuldeep Singh, Rajendra K Baharia, Mradul Mohan, Anup R Anvikar, Praveen K. Bharti, Age-specific malaria vulnerability and transmission reservoir among children, *Global Pediatrics*, 2023; 6, 100085.<https://doi.org/10.1016/j.gpeds.2023.100085>.
18. Arikawa, S., Tchankoni, M.K., Gbeasor-Komlanvi, F.A. *et al.* Prevalence and risk factors associated with malaria infection in children under two years of age in southern Togo prior to perennial malaria chemoprevention implementation. *Malar J* **22**,357(2023).<https://doi.org/10.1186/s12936-023-04793-y>
19. Debash H, Bisetegn H, Ebrahim H, Feleke DG, Gedefie A, Tilahun M, et al. (2022) Prevalence and associated risk factors of malaria among febrile under-five children visiting health facilities in Ziquala district, Northeast Ethiopia: A multicenter cross-sectional study. *PLoS ONE* 17(10): e0276899.<https://doi.org/10.1371/journal.pone.0276899>
20. Obasohan PE, Walters SJ, Jacques R, Khatab K. Individual and Contextual Factors Associated with Malaria among Children 6–59 Months in Nigeria: A Multilevel Mixed Effect Logistic Model Approach. *International Journal of Environmental Research and Public Health*. 2021;18(21):11234. <https://doi.org/10.3390/ijerph182111234>
21. Carter, R., Karunaweera, N.D. The role of improved housing and living environments in malaria control and elimination. *Malar J* **19**,385(2020).<https://doi.org/10.1186/s12936-020-03450-y>
22. Ugwu CLJ, Zewotir T. Evaluating the Effects of Climate and Environmental Factors on Under-5 Children Malaria Spatial Distribution Using Generalized Additive Models (GAMs). *J Epidemiol Glob Health*. 2020 Dec;10(4):304–314. doi: 10.2991/jegh.k.200814.001. Epub 2020 Aug 21. PMID: 33009733; PMCID: PMC7758859.
23. Gbenga Adegbite, Sunday Edeki, Itunuoluwa Isewon, Jerry Emmanuel, Titilope Dokunmu, Solomon Rotimi, Jelili Oyelade, Ezekiel Adebisi, Mathematical modeling of malaria transmission dynamics in humans with mobility and control states, *Infectious Disease Modelling*, 2023; 8(4): 10151031.<https://doi.org/10.1016/j.idm.2023.08.005>.
24. Fall P, Diouf I, Deme A, Diouf S, Sene D, Sultan B, Janicot S. Enhancing Understanding of the Impact of Climate Change on Malaria in West Africa Using the Vector-Borne Disease Community Model of the International Center for Theoretical Physics (VECTRI) and the Bias-Corrected Phase 6 Coupled Model Intercomparison Project Data (CMIP6). *Microbiology Research*. 2023; 14(4):2148–2180. <https://doi.org/10.3390/microbiolres14040145>
25. Alum EU, Ugwu OPC, Egba SI, Uti DE, Alum BN. (2024). Climate Variability and

- Malaria Transmission: Unraveling the Complex Relationship. *INOSR Scientific Research* 11(2):16-22. <https://doi.org/10.59298/INOSRSR/2024/1.1.21622>
26. Lena Fischer, Nejla Gültekin, Marisa B. Kaelin, Jan Fehr, Patricia Schlagenhauf, Rising temperature and its impact on receptivity to malaria transmission in Europe: A systematic review, *Travel Medicine and Infectious Disease*, 2020; 36, 101815. <https://doi.org/10.1016/j.tmaid.2020.101815>
27. Chiziba C, Mercer LD, Diallo O, Bertozzi-Villa A, Weiss DJ, Gerardin J, Ozodiegwu ID. Socioeconomic, Demographic, and Environmental Factors May Inform Malaria Intervention Prioritization in Urban Nigeria. *International Journal of Environmental Research and Public Health*. 2024; 21(1):78. <https://doi.org/10.3390/ijerph21010078>
28. Emmanuel Ifeanyi Obeagu, Getrude Uzoma Obeagu, Simeon Ikechukwu Egba and Obioma Raluchukwu Emeka Obi (2023) Combatting Anaemia in Paediatric Malaria: Effective management strategies *Int. J. Curr. Res. Med. Sci.* (2023). 9(11): 1-7
29. Mohammadkhani, M., Khanjani, N., Bakhtiari, B., Tabatabai, S. M., & Sheikhzadeh, K. (2019). The Relation Between Climatic Factors and Malaria Incidence in Sistan and Baluchestan, Iran. *SageOpen*, 9(3). <https://doi.org/10.1177/2158244019864205>
30. Emmanuel Ikechukwu Nnamonu., Ogonna Christiana Ani., Felix Joel Ugwu., Simeon Ikechukwu Egba., Ifeanyi Oscar Aguzie., Obiageli Panthe Okeke., Christian Enyi Dialoke., Lilian Obinna Asogwa and Solomon Ikechukwu Odo (2020) Malaria Prevalence in Rice Farm Settlements South East Nigeria. *IJTDH*, 2020; 41(9): 64-74
31. Abdulaziz Y.A. Mukhtar, Justin B. Munyakazi, Rachid Ouifki, Assessing the role of climate factors on malaria transmission dynamics in South Sudan, *Mathematical Biosciences*, 2019; 310: 13-23. <https://doi.org/10.1016/j.mbs.2019.01.002>
32. Dabaro, D., Birhanu, Z., Negash, A. *et al.* Effects of rainfall, temperature and topography on malaria incidence in elimination targeted district of Ethiopia. *Malar J* 20, 104 (2021). <https://doi.org/10.1186/s12936-021-03641-1>
33. Castro MC, Tsuruta A, Kanamori S, Kannady K, Mkude S. Community-based environmental management for malaria control: evidence from a small-scale intervention in Dar es Salaam, Tanzania. *Malar J*. 2009 Apr 8; 8:57. doi: 10.1186/1475-2875-8-57. PMID: 19356246; PMCID: PMC2683857.
34. Haq IU, Mehmood Z, Khan GA, Kainat B, Ahmed B, Shah J, Sami A, Nazar MS, Xu J and Xiang H (2024) Modeling the effect of climatic conditions and topography on malaria incidence using Poisson regression: a Retrospective study in Bannu, Khyber Pakhtunkhwa, Pakistan. *Front. Microbiol.* 14:1303087. doi: 10.3389/fmicb.2023.1303087
35. Agyemang-Badu SY, Awuah E, Oduro-Kwarteng S, Dzamesi JYW, Azure SA. Assessment of the Knowledge, Prevalence, and Control Strategies of Malaria Among Households in Sunyani Municipality, Bono Region, Ghana. *Environ Health Insights*. 2023 May 30; 17:11786302231176922. doi: 10.1177/11786302231176922. PMID: 37273838; PMCID: PMC10236243.
36. Bofu, R.M., Santos, E.M., Msugupakulya, B.J. *et al.* The needs and opportunities for housing improvement for malaria control in southern Tanzania. *Malar J* 22, 69 (2023). <https://doi.org/10.1186/s12936-023-04499-1>
37. Samsudin NA, Othman H, Siau CS, Zaini ZI. Exploring community needs in combating aedes mosquitoes and dengue fever: a study with urban community in the recurrent hotspot area. *BMC Public Health*. 2024 Jun 20; 24(1):1651. doi: 10.1186/s12889-024-18965-1. PMID: 38902692; PMCID: PMC11191142.

CITE AS: Katu Amina H. (2024). Socioeconomic and Environmental Risk Factors for Malaria in Young Children: A Review. *INOSR Experimental Sciences* 13(2):44-48. <https://doi.org/10.59298/INOSRES/2024/1324448.000>